

EFFICACY AND SAFETY OF DUAL-OPIOID TREATMENT WITH MORPHINE PLUS OXYCODONE FOR ACUTE POSTOPERATIVE PAIN

ROBIN KELEN, RN, BA, CNN¹; TONYA BRUNETTA, BS²

¹QRXPHARMA, INC., BEDMINSTER, NJ; ²INC RESEARCH, INC., AUSTIN, TX

INTRODUCTION

- Acute postoperative pain is frequently managed inadequately; if such pain persists, it can lead to a greater risk of developing chronic pain^{1,2}
- Opioids are the most commonly used drugs for acute postoperative pain³
 - At least 3 opioid receptors have been identified: mu (μ), kappa (κ), and delta (δ)
 - Opioid receptors are located in the brain and in various organs
 - Opioids interact differently with various receptors to produce diverse activity profiles
- Opioids, while effective in providing analgesia for most patients, are frequently accompanied by adverse events (AEs), such as nausea, vomiting, and dizziness, that limit their efficacy or cause early discontinuation
- Q8003 is a dual-opioid formulation that provides a fixed ratio (3:2) of morphine and oxycodone in an oral, immediate-release (IR) capsule for the management of postoperative pain
- The combination of morphine and oxycodone is believed to potentiate analgesia as a result of the interaction of these opioids on a broader spectrum of opioid subreceptors, including but not limited to μ and κ receptors^{4,7}
- Enhanced efficacy can result in the need to administer lower doses, leading to improved patient acceptance and response as well as the possibility for decreased side effects

OBJECTIVE

To determine the dose-responsive analgesic effects and safety of Q8003 in patients with moderate to severe pain following unilateral bunionectomy surgery

METHODS


Study design

- Randomized, double-blind, placebo-controlled, ascending-cohort, dose-response, 48-hour inpatient study following unilateral bunionectomy surgery
 - Q8003 was administered in 4 ascending cohorts: 3/2 mg, 6/4 mg, 12/8 mg, and 18/12 mg vs placebo
 - After completion of each dosing group, enrollment in the next higher dosing group began if not more than 10 Q8003-treated patients in the previous dosing group experienced intolerable opioid-related AEs
 - Q8003 was dosed prn with a required interdose interval of at least 1-2 hours to ensure patient safety (maximum of 5 doses in 12 hours for the 2 higher doses)
 - Rescue medication consisted of ibuprofen 600 mg
- Analysis of covariance (ANCOVA) was used to compare individual dosing cohorts vs placebo
- Time to perceptible analgesia was analyzed using Kaplan-Meier estimates

Patients


- Men and women ≥ 18 years of age with a body mass index (BMI) < 32 kg/m² who met the American Society of Anesthesiologists (ASA) Class I or II patient criteria, and who were scheduled for unilateral uncomplicated bunionectomy surgery, were eligible for the study
 - To continue in the study following surgery, patients were required to have:
 - Pain score of 2 or 3 on a 4-point Likert pain rating scale (moderate or severe intensity) (Figure 1)

FIGURE 1. LIKERT PAIN RATING SCALE



- Pain score of ≥ 4 on an 11-point Numerical Pain Rating Scale (NPRS; 0=no pain and 10=worst pain imaginable) (Figure 2)

FIGURE 2. NUMERICAL PAIN RATING SCALE



- Pulse oximetry score (SpO₂) of $\geq 95\%$ oxygen saturation
- Respiration rate of ≥ 12 breaths/minute

- Patients were excluded if they had significant opioid use in the prior year, a history of opioid intolerance, a history of drug abuse, or evidence of current alcohol abuse

Efficacy measures

- Primary efficacy measure
 - Mean sum of pain intensity differences (PIDs) by dose group during the 48-hour treatment period (SPID48)
 - SPID is an outcome measure that summarizes treatment response (ie, decrease in pain intensity) over a clinically relevant period
 - In this study, pain intensity was measured by the NPRS (Figure 2)
 - For the first dose of study medication, PIDs were calculated as the difference in the patient's PI score at baseline and the PI score at each scheduled post-treatment timepoint (15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, and 8 hours, as well as at the time of administration of rescue medication)
 - Following the second and all subsequent doses of study medication, PIDs were calculated as the difference between the PI score at the time of each dose and the PI score 1 hour after the dose

- Secondary efficacy measures
 - Global patient satisfaction using a 5-point scale (0=not effective to 4=excellent) recorded 48 hours after the initial dose
 - Onset of analgesia following first dose of study medication using the two stopwatch method
 - Mean morphine and oxycodone doses and mean interdose interval
 - Time to first dose of ibuprofen following first dose of study medication and mean number of ibuprofen doses
 - Number of patient withdrawals for lack of efficacy

Safety assessments


- Spontaneously reported AEs
- SpO₂ and respiration rate
- Opioid-Related Symptom Distress Scale patient checklist⁸
 - Rates occurrence of pre-defined opioid-related AEs
 - Completed 24 and 48 hours after initial dose of study medication

RESULTS

Patients

- Of 263 patients enrolled, 256 were randomly assigned to treatment and received ≥ 1 dose of study medication (Figure 3)

FIGURE 3. PATIENT DISPOSITION



- Patient demographics and baseline pain characteristics are presented in Table 1

TABLE 1. PATIENT DEMOGRAPHICS AND BASELINE PAIN CHARACTERISTICS

Characteristic	Placebo (n=60)	3/2 mg (n=48)	6/4 mg (n=49)	12/8 mg (n=50)	18/12 mg (n=49)
Age, y, mean (SD)	46.7 (14.5)	43.9 (14.6)	42.3 (13.6)	44.5 (13.1)	43.5 (12.8)
Gender, n (%)					
Men	14 (23)	13 (27)	5 (10)	8 (16)	9 (18)
Women	46 (77)	35 (73)	44 (90)	42 (84)	40 (82)
Race, n (%)					
White	44 (73)	34 (71)	38 (78)	31 (62)	32 (65)
Black	10 (17)	10 (21)	5 (10)	9 (18)	12 (25)
Asian	4 (7)	1 (2)	1 (2)	4 (8)	3 (6)
Hispanic	1 (2)	3 (6)	4 (8)	6 (12)	2 (4)
Other	1 (2)	0	1 (2)	0	0
BMI, kg/m ² , mean (SD)	26.4 (4.1)	25.6 (3.5)	25.0 (3.7)	25.0 (4.0)	25.4 (3.6)
NPRS, mean	6.6	6.2	6.7	6.6	6.8
Patients with moderate pain ^a , n (%)	46 (77)	39 (81)	39 (80)	40 (80)	34 (69)

^aRemainder of patients had severe pain at baseline. BMI=body mass index. NPRS=Numerical Pain Rating Scale (0=no pain; 10=worst pain imaginable).

Extent of exposure

- The mean dose of morphine/oxycodone per 6-hour period (mg/6h) and the mean interdose interval over the 48-hour study period are presented in Table 2
- The longest average interdose interval was observed in the 12/8-mg group (6.8 hours)


TABLE 2. STUDY MEDICATION DOSING

Dosing	Placebo (n=60)	3/2 mg (n=48)	6/4 mg (n=49)	12/8 mg (n=50)	18/12 mg (n=49)
Mean dose of morphine/oxycodone, mg/6h	0/0	6/4	9.8/6.5	11/7.3	15/10
Mean interdose interval, h	4.2	2.9	4.1	6.8	6.6

Efficacy

- SPID48 was significantly greater with each Q8003 dose compared with placebo (Figure 4)


FIGURE 4. MEAN SUM OF PAIN INTENSITY DIFFERENCES DURING THE 48-HOUR TREATMENT PERIOD (SPID48)



*P=0.0015 vs placebo, SD \pm 98; †P=0.0016 vs placebo, SD \pm 111; ‡P=0.0001 vs placebo, SD \pm 75; §P=0.0001 vs placebo, SD \pm 75

- The percentage of patients reporting "good" or "excellent" improvement was significantly greater with each dose of Q8003 compared with placebo (Figure 5)
- Patient satisfaction was greatest with the 12/8-mg dose (58.0% good or excellent, P<0.0001 vs placebo)

FIGURE 5. GLOBAL PATIENT SATISFACTION WITH Q8003 VS PLACEBO



- Following the first dose of study medication, patients who received Q8003 had a shorter time to confirmed perceptible analgesia (range across dose groups, 0.67 to 1.93 hours) compared with patients receiving placebo (4.4 hours)
 - The shortest time to confirmed perceptible analgesia was observed in the 12/8-mg group compared with the 3/2-mg, 6/4-mg, and 18/12-mg groups (0.67 hours vs 1.93, 1.04, and 0.73 hours, respectively)
- Following the first dose of study medication, patients who received Q8003 waited longer before taking ibuprofen (range across dose groups, 3.2 to 7.9 hours) compared with patients receiving placebo (2.9 hours)
- Patients receiving Q8003 required fewer doses of ibuprofen (mean, 1.9 to 2.3 doses/24h, range across dose groups) compared with patients receiving placebo (mean, 3.3 doses/24h)
- Fewer patients receiving Q8003 withdrew from the study for lack of efficacy (range across dose groups, 1 to 6 patients) compared with patients receiving placebo (15 patients)

Safety and tolerability

- The rate of discontinuation due to AEs was low in all treatment groups
 - Placebo, 2% (1 patient, hypotension); 3/2 mg, 2% (1 patient, nausea); 6/4 mg, 2% (1 patient, nausea); 12/8 mg, 8% (1 patient each: nausea, vomiting, decreased SpO₂, and hypotension); and 18/12 mg, 14% (2 patients each: nausea and vomiting; 1 patient each: hypoxia, pruritus, and hypotension)
- The most commonly reported AEs were those associated with opioid use (Table 3)
 - The majority of AEs were mild to moderate in intensity and the incidence decreased over time

TABLE 3. ADVERSE EVENTS OCCURRING IN $\geq 5\%$ OF PATIENTS BY DOSE GROUP AND ACTUAL DOSE (MEAN mg/6h)

Adverse event, n (%)	Placebo (0/0) (n=60)	3/2 mg (6/4) (n=48)	6/4 mg (9.8/6.5) (n=49)	12/8 mg (11/7.3) (n=50)	18/12 mg (15/10) (n=49)
Nausea	5 (8)	18 (38)	28 (57)	28 (56)	32 (65)
Vomiting	1 (2)	11 (23)	18 (37)	16 (32)	25 (51)
Dizziness	4 (7)	4 (8)	12 (25)	10 (20)	12 (25)
Headache	5 (8)	5 (10)	8 (16)	10 (20)	8 (16)
Pruritus/Generalized pruritus	0	3 (6)	10 (20)	9 (18)	9 (18)
SpO ₂ decrease	1 (2)	0	5 (10)	3 (6)	5 (10)
Somnolence	0	1 (2)	2 (4)	1 (2)	4 (8)
Constipation	1 (2)	3 (6)	3 (6)	6 (12)	2 (4)
Dry mouth	2 (3)	3 (6)	2 (4)	3 (6)	2 (4)

- In general, the percentages of patient-reported AEs on the Opioid-Related Symptom Distress Scale were higher than corresponding patient- and investigator-reported AEs; selected AEs for the entire treatment period are presented below:
 - Nausea was reported by 10% of patients receiving placebo and by 52%, 76%, 64%, and 78% of patients in the 3/2-mg, 6/4-mg, 12/8-mg, and 18/12-mg groups, respectively
 - Vomiting was reported by 0% of patients receiving placebo and by 23%, 37%, 40%, and 51% of patients in the 3/2-mg, 6/4-mg, 12/8-mg, and 18/12-mg groups, respectively
- Overall, 3 patients experienced serious AEs
 - 1 patient in the 18/12-mg group had hypotension that resulted in discontinuation of study medication
 - 2 patients in the placebo group had events not related to study medication (1, hand injury; 1, deep vein thrombosis)
- Desaturations were transient, infrequent, responded quickly to supplemental oxygen, and were more frequent in the 18/12-mg dose group than in other groups; none of the patients receiving Q8003 had a respiration rate of < 10 breaths/minute
- Sedation was seen in 1 patient in the 18/12-mg group; somnolence ranged from 2% to 8% across the Q8003 groups
- Euphoria was not observed

CONCLUSIONS

- Q8003 was associated with dose-related reductions in acute postoperative pain; even the lowest-dose group (mean 6/4 mg/6h) showed analgesic efficacy superior to placebo
- Q8003 was well tolerated, with few serious AEs or discontinuations, minimal changes in respiration and blood oxygenation, few occurrences of somnolence, and no occurrences of euphoria
- The low levels of somnolence and respiratory depression warrant further study
- The 12/8-mg dose appeared to provide the optimal combination of efficacy and tolerability; future studies will examine Q8003 with the dosing schedule of every 4 to 6 hours

REFERENCES

- Apfelbaum JL et al. *Anesth Analg*. 2003;97:534-40.
- Kehlet H et al. *Lancet*. 2006;367:1618-25.
- Krenzschick DA et al. *J Perianesth Nurs*. 2008;23:S28-42.
- Nielsen CK et al. *Pain*. 2007;132:289-300.
- Ross FB et al. *Pain*. 2000;84:421-8.
- Lauretti GR et al. *Br J Cancer*. 2003;89:2027-30.
- Blumenthal S et al. *Anesth Analg*. 2007;105:233-7.
- Gan TJ et al. *Acta Anaesthesiol Scand*. 2004;48:1194-1207.

Sponsored by QRxPharma, Bedminster, NJ, USA

Presented at the 2008 Annual Meeting of the American College of Nurse Practitioners, Nashville, Tennessee, October 29-November 2, 2008

EFFICACY AND SAFETY OF DUAL-OPIOID TREATMENT WITH MORPHINE PLUS OXYCODONE FOR ACUTE POSTOPERATIVE PAIN

ROBIN KELEN, RN, BA, CNN¹; TONYA BRUNETTA, BS²

¹QRXPHARMA, INC., BEDMINSTER, NJ; ²INC RESEARCH, INC., AUSTIN, TX

INTRODUCTION

- Acute postoperative pain is frequently managed inadequately; if such pain persists, it can lead to a greater risk of developing chronic pain^{1,2}
- Opioids are the most commonly used drugs for acute postoperative pain³
 - At least 3 opioid receptors have been identified: mu (μ), kappa (κ), and delta (δ)
 - Opioid receptors are located in the brain and in various organs
 - Opioids interact differently with various receptors to produce diverse activity profiles
- Opioids, while effective in providing analgesia for most patients, are frequently accompanied by adverse events (AEs), such as nausea, vomiting, and dizziness, that limit their efficacy or cause early discontinuation
- Q8003 is a dual-opioid formulation that provides a fixed ratio (3:2) of morphine and oxycodone in an oral, immediate-release (IR) capsule for the management of postoperative pain
- The combination of morphine and oxycodone is believed to potentiate analgesia as a result of the interaction of these opioids on a broader spectrum of opioid subreceptors, including but not limited to μ and κ receptors⁴⁻⁷
- Enhanced efficacy can result in the need to administer lower doses, leading to improved patient acceptance and response as well as the possibility for decreased side effects

OBJECTIVE

To determine the dose-responsive analgesic effects and safety of Q8003 in patients with moderate to severe pain following unilateral bunionectomy surgery

METHODS

Study design

- Randomized, double-blind, placebo-controlled, ascending-cohort, dose-response, 48-hour inpatient study following unilateral bunionectomy surgery
 - Q8003 was administered in 4 ascending cohorts: 3/2 mg, 6/4 mg, 12/8 mg, and 18/12 mg vs placebo
 - After completion of each dosing group, enrollment in the next higher dosing group began if not more than 10 Q8003-treated patients in the previous dosing group experienced intolerable opioid-related AEs
 - Q8003 was dosed prn with a required interdose interval of at least 1-2 hours to ensure patient safety (maximum of 5 doses in 12 hours for the 2 higher doses)
 - Rescue medication consisted of ibuprofen 600 mg
- Analysis of covariance (ANCOVA) was used to compare individual dosing cohorts vs placebo
- Time to perceptible analgesia was analyzed using Kaplan-Meier estimates

Patients

- Men and women ≥ 18 years of age with a body mass index (BMI) < 32 kg/m² who met the American Society of Anesthesiologists (ASA) Class I or II patient criteria, and who were scheduled for unilateral uncomplicated bunionectomy surgery, were eligible for the study
 - To continue in the study following surgery, patients were required to have:
 - Pain score of 2 or 3 on a 4-point Likert pain rating scale (moderate or severe intensity) (Figure 1)

FIGURE 1. LIKERT PAIN RATING SCALE

Please circle the one number that best describes how much pain you have <u>right now</u> :			
0	1	2	3
No Pain	Mild Pain	Moderate Pain	Severe Pain

- Pain score of ≥ 4 on an 11-point Numerical Pain Rating Scale (NPRS; 0=no pain and 10=worst pain imaginable) (Figure 2)

FIGURE 2. NUMERICAL PAIN RATING SCALE

Please circle the one number that best describes how much pain you have <u>right now</u> :										
0	1	2	3	4	5	6	7	8	9	10
No Pain										Worst pain imaginable

- Pulse oximetry score (SpO₂) of $\geq 95\%$ oxygen saturation
- Respiration rate of ≥ 12 breaths/minute
- Patients were excluded if they had significant opioid use in the prior year, a history of opioid intolerance, a history of drug abuse, or evidence of current alcohol abuse

Efficacy measures

- Primary efficacy measure
 - Mean sum of pain intensity differences (PIDs) by dose group during the 48-hour treatment period (SPID48)
 - SPID is an outcome measure that summarizes treatment response (ie, decrease in pain intensity) over a clinically relevant period
 - In this study, pain intensity was measured by the NPRS (Figure 2)
 - For the first dose of study medication, PIDs were calculated as the difference in the patient's PI score at baseline and the PI score at each scheduled post-treatment timepoint (15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, and 8 hours, as well as at the time of administration of rescue medication)
 - Following the second and all subsequent doses of study medication, PIDs were calculated as the difference between the PI score at the time of each dose and the PI score 1 hour after the dose


- Secondary efficacy measures
 - Global patient satisfaction using a 5-point scale (0=not effective to 4=excellent) recorded 48 hours after the initial dose
 - Onset of analgesia following first dose of study medication using the two stopwatch method
 - Mean morphine and oxycodone doses and mean interdose interval
 - Time to first dose of ibuprofen following first dose of study medication and mean number of ibuprofen doses
 - Number of patient withdrawals for lack of efficacy
- Safety assessments
 - Spontaneously reported AEs
 - SpO₂ and respiration rate
 - Opioid-Related Symptom Distress Scale patient checklist⁶
 - Rates occurrence of pre-defined opioid-related AEs
 - Completed 24 and 48 hours after initial dose of study medication

RESULTS

Patients

- Of 263 patients enrolled, 256 were randomly assigned to treatment and received ≥1 dose of study medication (Figure 3)

FIGURE 3. PATIENT DISPOSITION



- Patient demographics and baseline pain characteristics are presented in Table 1

TABLE 1. PATIENT DEMOGRAPHICS AND BASELINE PAIN CHARACTERISTICS

Characteristic	Placebo (n=60)	3/2 mg (n=48)	6/4 mg (n=49)	12/8 mg (n=50)	18/12 mg (n=49)
Age, y, mean (SD)	46.7 (14.5)	43.9 (14.6)	42.3 (13.6)	44.5 (13.1)	43.5 (12.8)
Gender, n (%)					
Men	14 (23)	13 (27)	5 (10)	8 (16)	9 (18)
Women	46 (77)	35 (73)	44 (90)	42 (84)	40 (82)
Race, n (%)					
White	44 (73)	34 (71)	38 (78)	31 (62)	32 (65)
Black	10 (17)	10 (21)	5 (10)	9 (18)	12 (25)
Asian	4 (7)	1 (2)	1 (2)	4 (8)	3 (6)
Hispanic	1 (2)	3 (6)	4 (8)	6 (12)	2 (4)
Other	1 (2)	0	1 (2)	0	0
BMI, kg/m ² , mean (SD)	26.4 (4.1)	25.6 (3.5)	25.0 (3.7)	25.0 (4.0)	25.4 (3.6)
NPRS, mean	6.6	6.2	6.7	6.6	6.8
Patients with moderate pain ^a , n (%)	46 (77)	39 (81)	39 (80)	40 (80)	34 (69)

^aRemainder of patients had severe pain at baseline.
 BMI=body mass index.
 NPRS=Numerical Pain Rating Scale (0=no pain; 10=worst pain imaginable).

Extent of exposure

- The mean dose of morphine/oxycodone per 6-hour period (mg/6h) and the mean interdose interval over the 48-hour study period are presented in **Table 2**
- The longest average interdose interval was observed in the 12/8-mg group (6.8 hours)


TABLE 2. STUDY MEDICATION DOSING

Dosing	Placebo (n=60)	3/2 mg (n=48)	6/4 mg (n=49)	12/8 mg (n=50)	18/12 mg (n=49)
Mean dose of morphine/oxycodone, mg/6h	0/0	6/4	9.8/6.5	11/7.3	15/10
Mean interdose interval, h	4.2	2.9	4.1	6.8	6.6

Efficacy

- SPID48 was significantly greater with each Q8003 dose compared with placebo (**Figure 4**)


FIGURE 4. MEAN SUM OF PAIN INTENSITY DIFFERENCES DURING THE 48-HOUR TREATMENT PERIOD (SPID48)



*P=0.0015 vs placebo, SD ± 98; †P=0.0016 vs placebo, SD ± 111; ‡P=0.0001 vs placebo, SD ± 75; §P<0.0001 vs placebo, SD ± 75

- The percentage of patients reporting “good” or “excellent” improvement was significantly greater with each dose of Q8003 compared with placebo (**Figure 5**)
- Patient satisfaction was greatest with the 12/8-mg dose (58.0% good or excellent, P<0.0001 vs placebo)

FIGURE 5. GLOBAL PATIENT SATISFACTION WITH Q8003 VS PLACEBO



- Following the first dose of study medication, patients who received Q8003 had a shorter time to confirmed perceptible analgesia (range across dose groups, 0.67 to 1.93 hours) compared with patients receiving placebo (4.4 hours)
 - The shortest time to confirmed perceptible analgesia was observed in the 12/8-mg group compared with the 3/2-mg, 6/4-mg, and 18/12-mg groups (0.67 hours vs 1.93, 1.04, and 0.73 hours, respectively)
- Following the first dose of study medication, patients who received Q8003 waited longer before taking ibuprofen (range across dose groups, 3.2 to 7.9 hours) compared with patients receiving placebo (2.9 hours)
- Patients receiving Q8003 required fewer doses of ibuprofen (mean, 1.9 to 2.3 doses/24h, range across dose groups) compared with patients receiving placebo (mean, 3.3 doses/24h)
- Fewer patients receiving Q8003 withdrew from the study for lack of efficacy (range across dose groups, 1 to 6 patients) compared with patients receiving placebo (15 patients)

Safety and tolerability

- The rate of discontinuation due to AEs was low in all treatment groups
 - Placebo, 2% (1 patient, hypotension); 3/2 mg, 2% (1 patient, nausea); 6/4 mg, 2% (1 patient, nausea); 12/8 mg, 8% (1 patient each: nausea, vomiting, decreased SpO₂, and hypotension); and 18/12 mg, 14% (2 patients each: nausea and vomiting; 1 patient each: hypoxia, pruritus, and hypotension)
- The most commonly reported AEs were those associated with opioid use (Table 3)
 - The majority of AEs were mild to moderate in intensity and the incidence decreased over time

TABLE 3. ADVERSE EVENTS OCCURRING IN ≥5% OF PATIENTS BY DOSE GROUP AND ACTUAL DOSE (MEAN mg/6h)

Adverse event, n (%)	Placebo (0/0) (n=60)	3/2 mg (6/4) (n=48)	6/4 mg (9.8/6.5) (n=49)	12/8 mg (11/7.3) (n=50)	18/12 mg (15/10) (n=49)
Nausea	5 (8)	18 (38)	28 (57)	28 (56)	32 (65)
Vomiting	1 (2)	11 (23)	18 (37)	16 (32)	25 (51)
Dizziness	4 (7)	4 (8)	12 (25)	10 (20)	12 (25)
Headache	5 (8)	5 (10)	8 (16)	10 (20)	8 (16)
Pruritus/Generalized pruritus	0	3 (6)	10 (20)	9 (18)	9 (18)
SpO ₂ decrease	1 (2)	0	5 (10)	3 (6)	5 (10)
Somnolence	0	1 (2)	2 (4)	1 (2)	4 (8)
Constipation	1 (2)	3 (6)	3 (6)	6 (12)	2 (4)
Dry mouth	2 (3)	3 (6)	2 (4)	3 (6)	2 (4)

- In general, the percentages of patient-reported AEs on the Opioid-Related Symptom Distress Scale were higher than corresponding patient- and investigator-reported AEs; selected AEs for the entire treatment period are presented below:
 - Nausea was reported by 10% of patients receiving placebo and by 52%, 76%, 64%, and 78% of patients in the 3/2-mg, 6/4-mg, 12/8-mg, and 18/12-mg groups, respectively
 - Vomiting was reported by 0% of patients receiving placebo and by 23%, 37%, 40%, and 51% of patients in the 3/2-mg, 6/4-mg, 12/8-mg, and 18/12-mg groups, respectively
- Overall, 3 patients experienced serious AEs
 - 1 patient in the 18/12-mg group had hypotension that resulted in discontinuation of study medication
 - 2 patients in the placebo group had events not related to study medication (1, hand injury; 1, deep vein thrombosis)
- Desaturations were transient, infrequent, responded quickly to supplemental oxygen, and were more frequent in the 18/12-mg dose group than in other groups; none of the patients receiving Q8003 had a respiration rate of <10 breaths/minute
- Sedation was seen in 1 patient in the 18/12-mg group; somnolence ranged from 2% to 8% across the Q8003 groups
- Euphoria was not observed

CONCLUSIONS

- Q8003 was associated with dose-related reductions in acute postoperative pain; even the lowest-dose group (mean 6/4 mg/6h) showed analgesic efficacy superior to placebo
- Q8003 was well tolerated, with few serious AEs or discontinuations, minimal changes in respiration and blood oxygenation, few occurrences of somnolence, and no occurrences of euphoria
- The low levels of somnolence and respiratory depression warrant further study
- The 12/8-mg dose appeared to provide the optimal combination of efficacy and tolerability; future studies will examine Q8003 with the dosing schedule of every 4 to 6 hours

REFERENCES

1. Apfelbaum JL et al. *Anesth Analg*. 2003;97:534-40.
2. Kehlet H et al. *Lancet*. 2006;367:1618-25.
3. Krenzschek DA et al. *J Perianesth Nurs*. 2008;23:S28-42.
4. Nielsen CK et al. *Pain*. 2007;132:289-300.
5. Ross FB et al. *Pain*. 2000;84:421-8.
6. Lauretti GR et al. *Br J Cancer*. 2003;89:2027-30.
7. Blumenthal S et al. *Anesth Analg*. 2007;105:233-7.
8. Gan TJ et al. *Acta Anaesthesiol Scand*. 2004;48:1194-1207.

Sponsored by QRxPharma, Bedminster, NJ, USA

Presented at the 2008 Annual Meeting of the American College of Nurse Practitioners, Nashville, Tennessee, October 29-November 2, 2008